

Subgradient and Subdifferential

Seminar

Optimization for ML. Faculty of Computer Science. HSE University

Main notions recap

i Main notions

For a domain set $E \in \mathbb{R}^n$ and a function $f : E \rightarrow \mathbb{R}$:

- A vector $g \in \mathbb{R}^n$ is called **subgradient** of the function f at $x \in E$ if $\forall y \in E$

$$f(y) \geq f(x) + g^T(y - x)$$

- A set $\partial f(x)$ is called **subdifferential** of the function f at $x \in E$ if:

$$\partial f(x) = \{g \in \mathbb{R}^n \mid f(y) \geq f(x) + g^T(y - x)\} \forall y \in E$$

- $f(\cdot)$ is called **subdifferentiable** at point $x \in E$ if $\partial f(x) \neq \emptyset$

Connection between subdifferentiation and convexity

💡 Connection between subdifferentiation and convexity

If $f : E \rightarrow \mathbb{R}$ is subdifferentiable on the **convex** subset $S \in E$ then f is convex on S .

- The inverse is generally incorrect
- There is no sense to derive the subgradient of nonconvex function.

Connection between subdifferentiation and differentiation

💡 Connection between subdifferentiation and differentiation

- 1) If $f : E \rightarrow \mathbb{R}$ is convex and differentiable at $x \in \text{int } E$ then $\partial f(x) = \{\Delta f(x)\}$
- 2) If $f : E \rightarrow \mathbb{R}$ is convex and for $x \in \text{int } E$ $\partial f(x) = \{s\}$ then f is differentiable at x and $\Delta f(x) = s$

- Derive the subdifferential of a differentiable function is overkill.

Problem 1

i Question

Find the subgradient of the function

$$f(x) = -\sqrt{x}$$

Subdifferentiation rules

1) $f : E \rightarrow \mathbb{R}, x \in E, c > 0$

$$\Rightarrow \partial(cf)(x) = c\partial f(x)$$

2) $f : F \rightarrow \mathbb{R}, g : G \rightarrow \mathbb{R}, x \in F \cap G$

$$\Rightarrow \partial(f + g)(x) \supseteq \partial f(x) + \partial g(x)$$

3) $T : V \rightarrow W = Ax + b, g : W \rightarrow \mathbb{R}, x_0 \in V$

$$\Rightarrow \partial(g \circ T)(x_0) \supseteq A^* \partial(g)(T(x_0))$$

4) $f(x) = \max(f_1(x), \dots, f_m(x)), I(x) = \{i \in 1 \dots m \mid f_i(x) = f(x)\}$

$$\Rightarrow \partial f(x) \supseteq \text{Conv} \left(\bigcup_{i \in I(x)} \partial f_i(x) \right)$$

💡 When is equality reached?

If abovementioned functions are convex and x is inner point then all inequalities turn into equalities.

Problem 2

Question

Find the subgradient of the function $f(x) + g(x)$ if

$$f(x) = -\sqrt{x} \text{ when } x \geq 0$$

$$g(x) = -\sqrt{-x} \text{ when } x \leq 0$$

Problem 3

i Question

- 1) Find the subgradient of the function $f(x) = \|Ax - b\|_1$;
- 2) For task $f(x) = \frac{1}{2}\|Ax - b\|_2^2 + \lambda\|x\|_1 \rightarrow \min_x$ say which lambdas lead to $x_{opt} = 0$

Problem 4

i Question

Check the differentiability of the function

$$f(A) = \sup_{\|x\|_2=1} x^T Ax, \text{ where } A \in \mathbb{S}^n, x \in \mathbb{R}^n$$